



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

ochraceous-brown on disk, lighter toward the margin which becomes crenulate and upturned in older specimens, slightly pruinose at first, rugulose and a little paler when dry: lamellæ slightly adnexed, not crowded, rather narrow, interspaced with anastomosing veins, much crisped, at first nearly white, then becoming ferruginous from the spores: stem 7-10^{cm} long, 0.1-0.3^{cm} thick, tapering from the somewhat bulbous base, yellowish-white, pruinose at base, hollow, fragile: spores 8-10 μ broad, 12-16 μ long.—In grass in dooryards and pastures, June and July.

The specific name is based on the peculiar character of the gills which are always crisped as soon as the pileus is expanded. Professor Charles H. Peck, to whom specimens of the above fungi were referred, and who very kindly reported on them, suggested that this might be a variety of *G. lateritia*, unless the peculiar character of the lamellæ proved to be constant. The fact that specimens possessing this character have been collected in the same localities during the past three seasons seems sufficient proof of the constancy of this character, and consequently, would indicate that this form is worthy of specific rank.—B. O. LONGYEAR, *Michigan Agricultural College*.

MEXICAN FUNGI. II.¹

THE following species of Ustilagineæ were collected by me in 1896 and 1898, and all, except the three marked with an asterisk, have been examined by P. Hennings, of Berlin.

USTILAGO RABENHORSTIANA P. Henn. On *Panicum filiforme*. Guadalajara, Oct. 12, 1896.

USTILAGO PAMPARUM Speg. On *Setaria*. City of Mexico, Oct. 2, 1896.

USTILAGO ULEI P. Henn. On *Chloris submutica*. City of Mexico, Oct. 2, 1896.

USTILAGO DIETELIANA P. Henn. On *Tripsacum dactyloides*. City of Mexico, Oct. 1, 1896.

USTILAGO HILARIÆ P. Henn. On *Hilaria cenchroides*. City of Mexico, Oct. 2, 1896.

USTILAGO ÆGOPOGINIS P. Henn. On *Ægopogon cenchroides* P. Henn. City of Mexico, Oct. 3, 1896.

¹For previous paper see BOT. GAZ. 24: 23. 1897.

USTILAGO BRUNKII Ell. and Gallw. In culms of *Andropogon perforatus*. Cardenas, Mexico, Oct. 22, 1898.

USTILAGO NUDA (Jens.) Kell. & Swingle. On *Hordeum vulgare*. Toluca, Sept. 19, 1898.

USTILAGO CENCHRI Lagerh. On *Cenchrus (tribuloides?)*. Cardenas, Oct. 22, 1898.

Ustilago Holwayana P. Henn., n. sp. Soris in spicis, eas destruentibus, longe cylindraceutis, primo membrana pallide cinerea velatis, atris; sporis subglobois vel ellipsoideis, subacutangulis, atrovioaceis, $13-16 \times 12-15 \mu$; episporo verrucoso, atro.—In spicis *Paspali velutini*. Patzcuaro, Michoacan, Oct. 19, 1898.

Ustilago Andropogonis-hirtifolii P. Henn., n. sp. Soris in paniculis, ovaria destruentibus, longe cylindraceutis, membrana cinnamomea vestitis, dein pulverulentis, atrovioaceis vel fusco-olivaceis, $9-14 \mu$; episporo granuloso verrucoso.—In paniculis *Andropogonis hirtifolii pubifloræ*. Patzcuaro, Michoacan, Oct. 20, 1898. Apparently quite distinct from all the species known on *Andropogon*.

Ustilago Panici-prolifera P. Henn., n. sp. Soris in paniculis, eo omnino destruentibus, membrana cinerea tectis, dein pulverulentis atrovioaceis; sporis subglobois, acutangulis, pallide olivaceo-fuscis, intus granulatis, $7-9 \mu$; episporo levi.—In paniculis *Panici prolifera acuminati*. City of Mexico, Oct. 10, 1898. This species differs from *U. Panici-miliacei* (Pers.) Wint. in its much smaller spores, and in having the sori covered with an ashy-gray membrane; and from *U. vesiculosa* P. Henn. and *U. pustulata* Tr. & Earle in its smooth lighter colored spores. The sori and general appearance are also different.

*USTILAGO PARLATOREI F. de Wald. On *Rumex Mexicanus* Meissn. Toluca, Mexico, Sept. 19, 1898.

*USTILAGO TRITICI (Pers.) Jens. On *Triticum vulgare*. City of Mexico, Oct. 11, 1898.

*USTILAGO ZEÆ (Beckm.) Ung. On *Zea Mays*. City of Mexico, Oct. 10, 1898.

CINTRACTIA AXICOLA Cornu. On *Fimbristylis*. Cuernavaca, Mexico, Sept. 26, 1898.

CINTRACTIA LEUCODERMA (Berk.) P. Henn. On *Rhynchospora*. Jalapa, Mexico, Oct. 3, 1898.—E. W. D. HOLWAY, *Decorah, Iowa*.